

11.9 and 11.11. RECOMMENDATION:

Should NURSES deliver the maintenance dose of magnesium sulphate to (a) prevent eclampsia and refer to a higher facility, and (b) treat eclampsia and refer to a higher facility?

Problem: Poor access to treatment for eclampsia
Option: Nurses delivering loading dose of magnesium sulphate to prevent and treat eclampsia
Comparison: Care delivered by other cadres or no care
Setting: Community/primary health care settings in LMICs with poor access to health professionals

Recommendation	<i>We recommend against the option</i>	<i>We suggest considering the option with targeted monitoring and evaluation</i>	<i>We recommend the option</i>
	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
We recommend against the use of nurses to deliver the maintenance dose of magnesium sulphate to prevent or treat eclampsia.			

Justification There is insufficient evidence on the effectiveness of nurses delivering a maintenance dose of magnesium sulphate to prevent or treat eclampsia and refer to a higher facility. In addition, the intervention is outside of their typical scope of practice and its acceptability is uncertain.

Implementation considerations - Not applicable

Monitoring and evaluation

Research priorities

11.9 and 11.11. EVIDENCE BASE:

Should NURSES deliver the maintenance dose of magnesium sulphate to (a) prevent eclampsia and refer to a higher facility, and (b) treat eclampsia and refer to a higher facility?

Problem: Poor access to treatment for eclampsia
Option: Nurses delivering loading dose of magnesium sulphate to prevent and treat eclampsia
Comparison: Care delivered by other cadres or no care
Setting: Community/primary health care settings in LMICs with poor access to health professionals

CRITERIA	JUDGEMENT	EVIDENCE	COMMENTS AND QUERIES															
BENEFITS & HARMS OF THE OPTIONS	<p>Are the anticipated desirable effects large?</p> <p>No <input type="checkbox"/> Probably no <input type="checkbox"/> Uncertain <input checked="" type="checkbox"/> Probably yes <input type="checkbox"/> Yes <input type="checkbox"/> Varies <input type="checkbox"/></p>	<p>One systematic review searched for studies that assessed the effects of nurse-led primary care compared to care that was given by primary care doctors (Laurant 2012). However, this review did not identify any studies that specifically assessed the effects of nurses delivering magnesium sulphate. We are therefore unable to draw any conclusions about the desirable or undesirable effects of this intervention.</p> <p>Indirect evidence: The same review identified a number of studies, mostly from high income settings, where nurses were compared to doctors for the delivery of other types of interventions. The review suggests that nurse care may improve several health outcomes while it may make no difference to other outcomes. However, the certainty of this evidence varies.</p> <table border="1"> <thead> <tr> <th>Outcomes</th> <th>Impacts</th> <th>Certainty of the anticipated effect</th> </tr> </thead> <tbody> <tr> <td>Patient health status</td> <td>For some outcomes, benefits in favour of nurses. For other outcomes, no differences between nurses and doctors</td> <td>Very low to moderate</td> </tr> <tr> <td>Patient mortality</td> <td>No differences between nurses and primary care doctors</td> <td>Moderate</td> </tr> <tr> <td>Process of care</td> <td>Mixed results: some studies showed differences between nurses and primary care doctors in process of care, e.g. nurses gave more advice to patients, while others showed no differences</td> <td>Very low to moderate</td> </tr> <tr> <td>Patient satisfaction and preferences</td> <td>Patients were significantly more satisfied with nurses compared with primary care doctors. Also, patients preferred significantly more often to see a nurse rather than a primary care doctor.</td> <td>Very low to moderate</td> </tr> </tbody> </table> <p>Annex: page 6 (Laurant 2012)</p>	Outcomes	Impacts	Certainty of the anticipated effect	Patient health status	For some outcomes, benefits in favour of nurses. For other outcomes, no differences between nurses and doctors	Very low to moderate	Patient mortality	No differences between nurses and primary care doctors	Moderate	Process of care	Mixed results: some studies showed differences between nurses and primary care doctors in process of care, e.g. nurses gave more advice to patients, while others showed no differences	Very low to moderate	Patient satisfaction and preferences	Patients were significantly more satisfied with nurses compared with primary care doctors. Also, patients preferred significantly more often to see a nurse rather than a primary care doctor.	Very low to moderate	<p>Note: A World Health Organisation guideline recommends that for settings where it is not possible to administer the full magnesium sulphate regimen, the use of magnesium sulphate loading dose, followed by immediate transfer to a higher-level health facility, is recommended for women with severe pre-eclampsia and eclampsia (very low quality evidence, weak recommendation) (WHO, 2011). The guideline makes no recommendation regarding (a) which cadre should deliver the loading or maintenance doses for preventing and treating eclampsia, and (b) what should be done when immediate transfer to a higher-level facility is not possible following the loading dose.</p>
	Outcomes		Impacts	Certainty of the anticipated effect														
	Patient health status		For some outcomes, benefits in favour of nurses. For other outcomes, no differences between nurses and doctors	Very low to moderate														
	Patient mortality		No differences between nurses and primary care doctors	Moderate														
Process of care	Mixed results: some studies showed differences between nurses and primary care doctors in process of care, e.g. nurses gave more advice to patients, while others showed no differences	Very low to moderate																
Patient satisfaction and preferences	Patients were significantly more satisfied with nurses compared with primary care doctors. Also, patients preferred significantly more often to see a nurse rather than a primary care doctor.	Very low to moderate																
<p>Are the anticipated undesirable effects small?</p> <p>No <input type="checkbox"/> Probably no <input type="checkbox"/> Uncertain <input checked="" type="checkbox"/> Probably yes <input type="checkbox"/> Yes <input type="checkbox"/> Varies <input type="checkbox"/></p>																		
<p>What is the certainty of the anticipated effects?</p> <p>Very low <input type="checkbox"/> Low <input type="checkbox"/> Moderate <input type="checkbox"/> High <input type="checkbox"/> No direct evidence <input checked="" type="checkbox"/> Varies <input type="checkbox"/></p>																		
<p>Are the desirable effects large relative to the undesirable effects?</p> <p>No <input type="checkbox"/> Probably no <input type="checkbox"/> Uncertain <input checked="" type="checkbox"/> Probably yes <input type="checkbox"/> Yes <input type="checkbox"/> Varies <input type="checkbox"/></p>																		
RESOURCE USE	<p>Are the resources required small?</p> <p>No <input type="checkbox"/> Probably no <input type="checkbox"/> Uncertain <input type="checkbox"/> Probably yes <input checked="" type="checkbox"/> Yes <input type="checkbox"/> Varies <input type="checkbox"/></p>	<p>Main resource requirements</p> <table border="1"> <thead> <tr> <th>Resource</th> <th>Settings in which nurses already provide other care</th> </tr> </thead> <tbody> <tr> <td>Training</td> <td>E.g. 2 weeks of practice-based training for nurses to diagnosis eclampsia and pre-eclampsia</td> </tr> <tr> <td>Supervision and monitoring</td> <td>Regular supervision by senior midwife or doctor</td> </tr> <tr> <td>Supplies</td> <td>Magnesium sulphate, calcium gluconate, IV equipment</td> </tr> <tr> <td>Referral</td> <td>Transportation to a centre where comprehensive emergency obstetric care (CeMOC) is available</td> </tr> </tbody> </table>	Resource	Settings in which nurses already provide other care	Training	E.g. 2 weeks of practice-based training for nurses to diagnosis eclampsia and pre-eclampsia	Supervision and monitoring	Regular supervision by senior midwife or doctor	Supplies	Magnesium sulphate, calcium gluconate, IV equipment	Referral	Transportation to a centre where comprehensive emergency obstetric care (CeMOC) is available						
Resource	Settings in which nurses already provide other care																	
Training	E.g. 2 weeks of practice-based training for nurses to diagnosis eclampsia and pre-eclampsia																	
Supervision and monitoring	Regular supervision by senior midwife or doctor																	
Supplies	Magnesium sulphate, calcium gluconate, IV equipment																	
Referral	Transportation to a centre where comprehensive emergency obstetric care (CeMOC) is available																	

	CRITERIA	JUDGEMENT	EVIDENCE	COMMENTS AND QUERIES
	<p>Is the incremental cost small relative to the benefits?</p>	<p>No <input type="checkbox"/> Probably no <input type="checkbox"/> Uncertain <input checked="" type="checkbox"/> Probably yes <input type="checkbox"/> Yes <input type="checkbox"/> Varies <input type="checkbox"/></p>	<p>Uncertain as there is no direct evidence on effectiveness. Indirect evidence from the review referred to above (Laurant 2012) suggests that, compared to doctor-led care:</p> <ul style="list-style-type: none"> • Overall, studies showed lower costs for nurse-led care • Consultation length was longer for nurses • For the frequency of consultations, results were mixed • For most studies there were no differences in the use of healthcare services and prescriptions 	
ACCEPTABILITY	<p>Is the option acceptable to most stakeholders?</p>	<p>No <input type="checkbox"/> Probably no <input type="checkbox"/> Uncertain <input type="checkbox"/> Probably yes <input checked="" type="checkbox"/> Yes <input type="checkbox"/> Varies <input type="checkbox"/></p>	<p>A systematic review of doctor-nurse substitution (Rashidian 2012) did not identify any studies that evaluated the acceptability of the loading dose of magnesium sulphate for eclampsia when delivered by nurses. We are therefore uncertain about the acceptability of this intervention to key stakeholders.</p> <p>For other maternal and child health interventions, the same review suggests that:</p> <ul style="list-style-type: none"> • Nurses may be motivated to offer advanced care by increased recognition and job satisfaction (moderate certainty evidence) • Recipients may regard nurses as more accessible and better at listening and caring than doctors (moderate certainty evidence). However, some recipients may have concerns about nurses' competence and willingness to provide high quality care compared to doctors (low certainty evidence). In addition, for tasks that are more "medical" in nature, recipients may prefer doctors over nurses (low certainty evidence) • Doctors were generally satisfied with the contribution of nurses to maternal and child health care, although some concerns were raised (low certainty evidence). Doctors may welcome the contribution of nurses where it reduces doctors' workloads (moderate certainty evidence). Doctor acceptance may be influenced by level of nurse experience (low certainty evidence). Doctors may be comfortable with nurse prescribing, believing that it improves continuity of care (low certainty evidence). However, an increase in nurse autonomy may negatively affect or produce negative reactions among other professions, including doctors and midwives, who for instance may be unwilling to relinquish final responsibility for patient care. A lack of clarity about nurse roles and responsibilities in relation to other health workers may also be a challenge (low certainty evidence) <p>Annex: page 43 (Rashidian 2012)</p>	
FEASIBILITY	<p>Is the option feasible to implement?</p>	<p>No <input type="checkbox"/> Probably no <input type="checkbox"/> Uncertain <input type="checkbox"/> Probably yes <input checked="" type="checkbox"/> Yes <input type="checkbox"/> Varies <input type="checkbox"/></p>	<p>The intervention requires relatively few supplies (Magnesium sulphate, calcium gluconate, IV equipment). In addition, it is simple to deliver and requires only a small amount of training.</p> <p>Regular supervision needs to be in place, and adequate referral to a higher level of care for further management may also be necessary. However, systematic review (Rashidian 2012) suggests that nurses may be unprepared or not adequately trained or supervised when they are given advanced and substitution roles (low certainty). In some settings, changes to norms or regulations may be needed to allow nurses to prescribe and deliver the loading dose of magnesium sulphate.</p> <p>Annex: page 43 (Rashidian 2012)</p>	